

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018697 A2

(51) International Patent Classification⁷: **C12Q**

(21) International Application Number:
PCT/US2003/025975

(22) International Filing Date: 19 August 2003 (19.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/166,356 21 August 2002 (21.08.2002) US

(71) Applicant: **BAYLOR COLLEGE OF MEDICINE**
[US/US]; One Baylor Plaza, Houston, TX 77030 (US).

(72) Inventors: **DRAGHIA-AKLI, Ruxandra**; 5215
Starkridge Drive, Houston, TX 77035 (US). **FIOROTTO,**
Marta, L.; 3706 Nottingham, Houston TX 77005 (US).
TAFFET, George; 4131 Turnberry Circle, Houston, TX
77025-1715 (US).

(74) Agents: **CHWANG, T., Ling** et al.; Jackson Walker L.L.P.,
2435 N. Central Expressway, Suite 600, Richardson, TX
75080 (US).


(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— without international search report and to be republished
upon receipt of that report

[Continued on next page]

(54) Title: PROTEASE RESISTANT TI-GROWTH HORMONE RELEASING HORMONE



(57) **Abstract:** One aspect of the current invention is a composition for a modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof. Another aspect of the current invention includes a nucleic acid molecule that encodes the modified GHRH or functional biological equivalent. The modified GHRH can be defined as a biologically active polypeptide that was engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to a wild-type GHRH ("wt-GHRH") polypeptide. Another aspect of the current invention includes a method for delivering the composition of this invention to a subject, wherein the modified GHRH increases the level of growth hormone ("GH") secretion in a subject. The preferred subject is a human or domesticated animal. Additionally, the modified GHRH composition is resistant to degradation when compared to the wt-GHRH.

WO 2004/018697 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

PROTEASE RESISTANT TI-GROWTH HORMONE RELEASING HORMONE RELATED APPLICATIONS

[0001] This application is a continuation-in-part of the U.S. Patent Application, Serial Number 09/624,268, entitled "Super-Active Porcine Growth Hormone Releasing Hormone Analog," filed on 7/24/2000. The 09/624,268 application claims priority to U.S. Provisional Patent Application, Serial Number 60/145,624, entitled "Super-Active Porcine Growth Hormone Releasing Hormone Analog," filed on 7/26/1999, the entire content of each of which is hereby incorporated by reference.

BACKGROUND

[0002] One aspect of the current invention is a composition for a modified growth hormone releasing hormone ("GHRH") molecule or functional biological equivalent thereof. The composition may also be a nucleic acid molecule that encodes the modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof. The modified GHRH can be defined as a biologically active polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to a wild-type GHRH ("wt-GHRH") polypeptide. One benefit of the claimed invention occurs when the modified molecule with the GHRH composition is delivered to a subject. The modified GHRH increases the level of growth hormone ("GH") secretion in a subject. Other benefits outlined in preferred embodiments include: increased insulin-like growth factor I ("IGF-I"), increased weight gain; increased bone density, increases body length, improved immune function and improved cardiac function in aging mammals. The modified GHRH composition of this invention is also resistant to degradation compared to the wt-GHRH.

[0003] Regulated expression of the growth hormone ("GH") pathway is essential for optimal linear growth, as well as homeostasis of carbohydrate, protein, and fat metabolism. GH synthesis and its pulsatile secretion from the anterior pituitary is stimulated by growth hormone releasing hormone ("GHRH") and inhibited by somatostatin, both hypothalamic hormones (Frohman et al., 1992). GH increases production of insulin-like growth factor-I (IGF-I) primarily in the liver, as well as other

target organs. IGF-I and GH feedback on the hypothalamus and pituitary to inhibit GHRH release and GH secretion. The endogenous rhythm of GH secretion becomes entrained to the imposed rhythm of exogenous GHRH (Caroni and Schneider, 1994).

[0004] Linear growth velocity and body composition respond to GH or GHRH replacement therapies in a broad spectrum of conditions, both in humans and in farm animals. The etiology of these conditions can vary significantly. In 50% of human GH deficiencies the GHRH-GH-IGF-I axis is functionally intact, but does not elicit the appropriate biological responses in its target tissues. Similar phenotypes are produced by genetic defects at different points along the GH axis (Parks et al., 1995), as well as in non-GH-deficient short stature. In the non-GH-deficiency causes of short stature, such as Turner syndrome (Butler et al., 1994), hypochondroplasia (Foncea et al., 1997), Crohn's disease (Parrizas and LeRoith, 1997), intrauterine growth retardation (Hoess and Abremski, 1985) or chronic renal insufficiency (Lowe, Jr. et al., 1989), GHRH or GH therapy can be effective in promoting linear growth (Gesundheit and Alexander, 1995). In the elderly, the GHRH-GH-IGF-I axis undergoes considerable decrement, with reduced GH secretion and IGF-I production associated with a loss of skeletal muscle mass (sarcopenia), osteoporosis, increased fat deposition and decreased lean body mass (Caroni and Schneider, 1994; Veldhuis et al., 1997). It has been demonstrated that the development of these changes can be offset by recombinant GH therapy. GH replacement therapy both in children and the elderly is widely used clinically. Current GH therapy has several shortcomings, however, including frequent subcutaneous or intravenous injections, insulin resistance and impaired glucose tolerance (Rabinovsky et al., 1992); children are also vulnerable to premature epiphyseal closure and slippage of the capital femoral epiphysis (Liu and LeRoith, 1999). A "slow-release" form of GH (Genentech), which requires injections every 14 days, perturbs the normal physiological pulsatile GH profile, and is also associated with frequent side effects.

[0005] In domestic livestock, GHRH and GH stimulate milk production, with an increase in feed to milk conversion, which additionally enhances growth, primarily by increasing lean body mass (Lapierre et al., 1991; van Rooij et al., 2000) with overall improvement in feed efficiency. Hot and chilled carcass weights are increased and

carcass lipid (percent of soft-tissue mass) is decrease by GHRH and GH (Etherton et al., 1986).

[0006] Administering novel GHRH analog proteins (U.S. Pat Nos. 5,847,066; 5,846,936; 5,792,747; 5,776,901; 5,696,089; 5,486,505; 5,137,872; 5,084,442, 5,036,045; 5,023,322; 4,839,344; 4,410,512, RE33,699), synthetic or naturally occurring peptide fragments of GHRH (U.S. Pat. Nos. 4,833,166; 4,228,158; 4,228,156; 4,226,857; 4,224,316; 4,223,021; 4,223,020; 4,223, 019) for the purpose of increasing release of growth hormone have been reported. GHRH analogs containing the following mutations have been reported (U.S. Patent No. 5,846,936): Tyr at position 1 to His; Ala at position 2 to Val, Leu, or others; Asn at position 8 to Gln, Ser, or Thr; Gly at position 15 to Ala or Leu; Met at position 27 to Nle or Leu; and Ser at position 28 to Asn. The GHRH analog is the subject of U.S. Patent Application Serial No. 09/624,268 ("the '268 patent"), which teaches application of a GHRH analog containing mutations that improve the ability to elicit the release of growth hormone. In addition, the '268 patent relates to the treatment of growth deficiencies; the improvement of growth performance; the stimulation of production of growth hormone in an animal at a greater level than that associated with normal growth; and the enhancement of growth utilizing the administration of growth hormone releasing hormone analog and is herein incorporated by reference.

[0007] Although GHRH protein therapy stimulates normal cyclical GH secretion with virtually no side effects (Corpas et al., 1993), the short half-life of the molecule *in vivo* requires frequent (one to three times per day) intravenous, subcutaneous or intranasal (at 300-fold higher dose) administrations. Thus, recombinant GHRH administration is not practical as a chronic therapy. However, extracranially secreted GHRH, as a mature or a truncated polypeptide, is often biologically active (Thorner et al., 1984) and a low level of serum GHRH (100pg/ml) stimulates GH secretion (Corpas et al., 1993). These characteristics make GHRH an excellent candidate for plasmid mediated supplementation of a gene product.

[0008] **Gene Delivery and *in vivo* Expression:** Recently, the delivery of specific genes to somatic tissue in a manner that can correct inborn or acquired deficiencies and imbalances was proved to be possible. Gene-based drug delivery offers a number of advantages over the administration of recombinant proteins. These advantages

include the conservation of native protein structure, improved biological activity, avoidance of systemic toxicities, and avoidance of infectious and toxic impurities. In addition, gene therapy allows for prolonged exposure to the protein in the therapeutic range, because the protein is synthesized and secreted continuously into the circulation.

[0009] The primary limitation of using recombinant protein is the limited availability of protein after each administration. Gene therapy using injectable DNA plasmid vectors overcomes this, because a single injection into the patient's skeletal muscle permits physiologic expression of the protein for extensive periods of time (WO 99/05300 and WO 01/06988). Injection of the vectors promotes the production of enzymes and hormones in animals in a manner that more closely mimics the natural process. Furthermore, among the non-viral techniques for gene transfer *in vivo*, the direct injection of plasmid DNA into muscle tissue is simple, inexpensive, and safe.

[0010] Direct plasmid DNA transfer is currently the basis of many emerging therapeutic strategies, as it avoids the potential problems associated with viral genes or lipid particles (Muramatsu et al., 1998). Skeletal muscle is a preferred target tissue, because the muscle fiber has a long life span and can be transduced by circular DNA plasmids. Skeletal muscle borne plasmids have been expressed efficiently over months or years in immunocompetent hosts (Davis et al., 1993; Tripathy et al., 1996). Plasmid DNA constructs are attractive candidate for direct therapy into the subjects skeletal muscle because they are well-defined entities, which are biochemically stable and have been used successfully for many years (Acsadi et al., 1991; Wolff et al., 1990). The relatively low expression levels, achieved after direct plasmid DNA injection are sometimes sufficient to prove bio-activity of secreted peptides (Danko and Wolff, 1994; Tsurumi et al., 1996).

[0011] Previously, we reported that in mice, human GHRH cDNA could be delivered to muscle by an injectable myogenic expression vector, where it transiently stimulated GH secretion over a period of two weeks (Draghia-Akli et al., 1997). We have then optimized this injectable vector system by incorporating a powerful synthetic muscle promoter (Li et al., 1999) coupled with a novel protease-resistant GHRH molecule with a substantially longer half-life and greater GH secretory activity (pSP-HV-GHRH) (Draghia-Akli et al., 1999). We improved vector delivery to skeletal muscle via a highly efficient electroporation technology (Wang et al., 1998). Using this combination of vector design and

electric pulses plasmid delivery method, we were able to show increased growth and favorably modified body composition in pigs (Draghia-Akli et al., 1999) and rodents (Draghia-Akli et al., 2002). The current invention describes a new protease resistant, super-active GHRH analog, called TI-GHRH.

SUMMARY

[0012] One aspect of the current invention is a composition for a modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof. A preferred embodiment of this invention includes a peptide with a general formula (-A₁-A₂-DAIFTNSYRKVL-A₃-QLSARKLLQDI-A₄-A₅-RQQGERNQQGA-OH), wherein A₁ is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H"); A₂ is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I"); A₃ is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G"); A₄ is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L"); A₅ is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N"). Other preferred embodiments include a modified GHRH or functional biological equivalent thereof as shown in Seq ID #3, Seq ID #4, and Seq ID #5. The modified GHRH can be defined as a biologically active polypeptide that was engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity compared to the wild-type GHRH ("wt-GHRH") polypeptide. One benefit of the claimed invention occurs when the modified GHRH composition is delivered to a subject. The modified GHRH increases the level of growth hormone ("GH") secretion in a subject. The preferred subject is a domesticated animal or human. Other benefits outlined in preferred embodiments include: increased insulin-like growth factor I ("IGF-I"), increased weight gain; increased bone density, increases body length, improved immune function and improved cardiac function in aging mammals. Additionally, the modified GHRH composition is resistant to degradation when compared to the wt-GHRH.

[0013] Another aspect of the current invention is a nucleic acid molecule (e.g. DNA or RNA) that encodes the modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof. In a preferred embodiment, nucleic acid molecule further comprises a synthetic mammalian expression plasmid with a synthetic or eukaryotic promoter; and a poly adenelation signal; a selectable marker gene promoter; a ribosomal binding site; and an origin of replication. The synthetic or eukaryotic promoter, the nucleic acid sequence encoding the modified GHRH or functional biological equivalent thereof, and the poly-adenylation signal comprise therapeutic elements of the synthetic mammalian expression plasmid. The therapeutic

elements are operatively linked and located in a first operatively-linked arrangement. Similarly, the selectable marker gene promoter, the ribosomal binding site, the selectable marker gene sequence, and the origin of replication comprise replication elements of the synthetic mammalian expression plasmid; the replication elements are operatively linked and located in a second operatively-linked arrangement. The first-operatively-linked arrangement and the second-operatively-linked arrangement comprise a circular structure of the synthetic mammalian expression plasmid. In a preferred embodiment, the synthetic mammalian expression plasmid is utilized for plasmid mediated gene supplementation. Other preferred embodiments of a nucleic acid molecule that encodes a modified GHRH comprise a 3' untranslated region ("UTR") encompassing a portion of a human growth hormone 3'UTR, and a modified myogenic promoter (e.g. pSPc5 12).

[0014] In preferred embodiments, the nucleic acid molecule that encodes the modified GHRH or functional biological equivalent thereof is combined with a transfection-facilitating polypeptide (e.g. poly-L-glutamate) for delivering the composition into a muscle cell of a subject. The encoded modified GHRH or functional biological equivalent thereof is expressed in a tissue specific manner in the subject.

BRIEF DESCRIPTION OF FIGURES

[0015] Figure 1 shows the amino acid sequence for porcine wild type growth hormone releasing hormone ("GHRH") and a protease resistant TI-GHRH analog with extended activity that can increase GH secretory activity and stability;

[0016] Figure 2 shows a the release of growth hormone ("GH") in porcine primary pituitary culture that were simulated by different GHRH species isolated from conditioned media of skeletal muscle cells transfected with myogenic expression vectors driving porcine GHRH analogs; the analogs are denoted as follows: porcine wild type GHRH (1-40)OH ("pwt"); pGHRH with amino acid substitutions of Gly15 to Ala, Met27 to Leu and Ser28 to Asn is represented by ("15/27/28"); the 15/27/28 construct plus the conversion of Ala2 to Ile2 is represented by ("TI-GHRH"); the 15/27/28 construct plus the conversion of Ala2 to Val2 is represented by ("TV-GHRH"); the 15/27/28 construct plus conversion of Tyr1 with His, and Ala2 with Val is represented by ("HV-GHRH"); a construct coding for *E.coli* beta-galactosidase ("βgal") is used as a negative control; and a positive control of recombinant human ("10ng GHRH");

[0017] Figure 3 shows the enhanced stability of the TI-GHRH compared with wild type porcine GHRH over a 6 hour incubation in plasma;

[0018] Figure 4 shows a schematic representation of three plasmid constructs: the porcine wild type ("pGHRH"); the TI-GHRH; and the β-galactosidase construct; all contain the SPc5-12 synthetic promoter and the 3' untranslated region ("UTR") of growth hormone ("GH"), kan (kanamycin resistance gene for bacterial selection), and neo (neomycin resistance gene for *in vivo* selection);

[0019] Figure 5 shows the relative levels of serum IGF-I concentration in pSP-GHRH injected mice versus placebo injected mice that were exposed to a single injection of any one of the super analog GHRH myogenic expression vectors;

[0020] Figure 6 shows the average body weight in mice after pSP-GHRH analogs were injected intra-muscularly compared to controls;

[0021] Figure 7 shows the average lean body mass increase in mice after pSP-GHRH analogs were injected intra-muscularly compared to controls;

[0022] Figure 8 shows the average bone area increase in mice after pSP-GHRH analogs were injected intra-muscularly compared to controls;

[0023] Figure 9 shows the average body length increase in mice after TI-GHRH analogs were injected intra-muscularly compared to controls;

[0024] Figure 10 shows the average tibia length in mice after TI-GHRH analogs were injected intra-muscularly compared to controls;

[0025] Figure 11 shows the average spleen weight in mice after TI-GHRH analogs were injected intra-muscularly compared to controls.

[0026] Figure 12 shows the E peak early filling velocity of mice heart after TI-GHRH analogs were injected intra-muscularly compared with controls. Heart rate and A peak velocity are shown as control measures.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0027] *Terms:*

[0028] The term “analog” as used herein includes any mutant of GHRH, or synthetic or naturally occurring peptide fragments of GHRH.

[0029] The term “codon” as used herein refers to any group of three consecutive nucleotide bases in a given messenger RNA molecule, or coding strand of DNA that specifies a particular amino-acid, or a starting or stopping signal for translation. The term codon also refers to base triplets in a DNA strand.

[0030] The term “coding region” as used herein refers to any portion of the DNA sequence that is transcribed into messenger RNA (mRNA) and then translated into a sequence of amino acids characteristic of a specific polypeptide.

[0031] The term “delivery” as used herein is defined as a means of introducing a material into a subject, a cell or any recipient, by means of chemical or biological process, injection, mixing, electroporation, sonoporation, or combination thereof, either without or under pressure.

[0032] The term “encoded GHRH” as used herein is a biologically active polypeptide.

[0033] The term “functional biological equivalent” of GHRH as used herein is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to the GHRH polypeptide.

[0034] The term “growth hormone” (“GH”) as used herein is defined as a hormone that relates to growth and acts as a chemical messenger to exert its action on a target cell.

[0035] The term “growth hormone releasing hormone” (“GHRH”) as used herein is defined as a hormone that facilitates or stimulates release of growth hormone, and to a lesser extent other pituitary hormones, such as prolactin.

[0036] The term “heterologous nucleic acid sequence” as used herein is defined as a DNA sequence consisting of differing regulatory and expression elements.

[0037] The term “modified GHRH” as used herein is a polypeptide that has been engineered to contain an amino acid sequence that is distinct from the wild-type GHRH polypeptide while simultaneously having similar or improved biological activity when compared to the wild-type GHRH polypeptide. The wild-type GHRH polypeptide is the naturally occurring species-specific GHRH polypeptide of a subject, a cell or any recipient of the modified GHRH.

[0038] The term “nucleic acid expression construct” as used herein refers to any type of genetic construct comprising a nucleic acid coding for a RNA capable of being transcribed. The transcribed RNA is then capable of being translated into a peptide, polypeptide, or protein. The term “expression vector” or “expression plasmid” can also be used interchangeably.

[0039] The term “subject” as used herein refers to any species of the animal kingdom. In preferred embodiments it refers more specifically to humans and domesticated animals.

[0040] The term “domesticated animal” as used herein refers to animals used for: pets (e.g. cats, dogs, etc.); work (e.g. horses, cows, etc.); food (chicken, fish, lambs, pigs, etc); and all others known in the art.

[0041] The term “operatively linked” as used herein refers to elements or structures in a nucleic acid sequence that are linked by operative ability and not physical location. The elements or structures are capable of, or characterized by accomplishing a desired operation. It is recognized by one of ordinary skill in the art that it is not necessary for elements or structures in a nucleic acid sequence to be in a tandem or adjacent order to be operatively linked.

[0042] The term “promoter” as used herein refers to a sequence of DNA that directs the transcription of a gene. A promoter may direct the transcription of a prokaryotic or eukaryotic gene. A promoter may be “inducible,” initiating transcription in response to an inducing agent or, in contrast, a promoter may be “constitutive,” whereby

an inducing agent does not regulate the rate of transcription. A promoter may be regulated in a tissue-specific or tissue-preferred manner, such that it is only active in transcribing the operable linked coding region in a specific tissue type or types.

[0043] The term "replication element" as used herein comprises nucleic acid sequences that will lead to replication of a plasmid in a specified host. One skilled in the art of molecular biology will recognize that the replication element may include, but is not limited to, a selectable marker gene promoter, a ribosomal binding site, a selectable marker gene sequence, and an origin of replication.

[0044] The term "therapeutic element" as used herein comprises nucleic acid sequences that will lead to an *in vivo* expression of an encoded gene product. One skilled in the art of molecular biology will recognize that the therapeutic element may include, but is not limited to a promoter sequence, a poly [A] sequence, or a 3' or 5' UTR.

[0045] The term "vector" as used herein refers to any vehicle that delivers a nucleic acid into a cell or organism. Examples include plasmid vectors, viral vectors, liposomes, or cationic lipids.

[0046] The standard one and three letter abbreviations for amino acids used herein are as follows: Alanine, A ala; Arginine, R, arg; Asparagine, N, asn; Aspartic acid, D, asp; Cysteine, C, cys; Glutamine, Q, gln; Glutamic acid, E, glu; Glycine, G, gly; Histidine, H, his; Isoleucine, I, ile; Leucine, L, leu; Lysine, K, lys; Methionine, M, met; Phenylalanine, F, phe; Proline, P, pro; Serine, S, ser; Threonine, T, thr; Tryptophan, W, trp; Tyrosine, Y, tyr; Valine, V, val.

[0047] In a preferred embodiment, the nucleic acid construct or vector of the present invention is a plasmid which comprises a synthetic myogenic (muscle-specific) promoter, a synthetic nucleotide sequence encoding a modified growth hormone releasing hormone (GHRH) or its analog, and a 3' untranslated region (3'UTR).

[0048] **Promoters and Enhancers.** A "promoter" is a control sequence that is a region of a nucleic acid sequence at which the initiation and rate of transcription are controlled. It may contain genetic elements where regulatory proteins and molecules may bind such as RNA polymerase and transcription factors. The phrases "operatively

positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and/or expression of that sequence. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0049] A promoter may be one of naturally-coding sequences located upstream of the coding segment and/or exon. Such a promoter can be referred to as "endogenous." Similarly, an enhancer may be one naturally associated with a nucleic acid sequence, located either downstream or upstream of that sequence. Alternatively, certain advantages will be gained by positioning the coding nucleic acid segment under the control of a recombinant or heterologous promoter, which refers to a promoter that is not normally associated with a nucleic acid sequence in its natural environment. A recombinant or heterologous enhancer refers also to an enhancer not normally associated with a nucleic acid sequence in its natural environment. Such promoters or enhancers may include promoters or enhancers of other genes, and promoters or enhancers isolated from any other prokaryotic, viral, or eukaryotic cell, and promoters or enhancers not "naturally occurring," *i.e.*, containing different elements of different transcriptional regulatory regions, and/or mutations that alter expression. In addition to producing nucleic acid sequences of promoters and enhancers synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including PCR™. Furthermore, it is contemplated the control sequences that direct transcription and/or expression of sequences within non-nuclear organelles such as mitochondria, chloroplasts, and the like, can be employed as well.

[0050] Naturally, it will be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the cell type, organelle, and organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression. The promoters employed may be constitutive, tissue-specific, inducible, and/or useful under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of

recombinant proteins and/or peptides. The promoter may be heterologous or endogenous. In a specific embodiment the promoter is a synthetic myogenic promoter.

[0051] The identity of tissue-specific promoters or elements, as well as assays to characterize their activity, is well known to those of skill in the art. Examples of such regions include the human LIMK2 gene, the somatostatin receptor 2 gene, murine epididymal retinoic acid-binding gene, human CD4, mouse alpha2 (XI) collagen, D1A dopamine receptor gene, insulin-like growth factor II, human platelet endothelial cell adhesion molecule-1.

[0052] Initiation Signals and Internal Ribosome Binding Sites. A specific initiation signal also may be required for efficient translation (synthesis of the encoded protein) of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements.

[0053] In certain embodiments of the invention, the use of internal ribosome entry sites ("IRES") elements are used to create multigene, or polycistronic messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap-dependent translation and begin translation at internal sites. IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described, as well as an IRES from a mammalian message. IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message.

[0054] **Multiple Cloning Sites.** Vectors can include a multiple cloning site ("MCS"), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector. "Restriction enzyme digestion" refers to catalytic cleavage of a nucleic acid molecule with an enzyme that functions only at specific locations in a nucleic acid molecule. Many of these restriction enzymes are commercially available. Use of such enzymes is widely understood by those of skill in the art. Frequently, a vector is linearized or fragmented using a restriction enzyme that cuts within the MCS to enable exogenous sequences to be ligated to the vector. "Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments, which may or may not be contiguous with each other. Techniques involving restriction enzymes and ligation reactions are well known to those of skill in the art of recombinant technology.

[0055] **Splicing Sites.** Most transcribed eukaryotic RNA molecules will undergo RNA splicing to remove introns from the primary transcripts. Vectors containing genomic eukaryotic sequences may require donor and/or acceptor splicing sites to ensure proper processing of the transcript for protein expression.

[0056] **Polyadenylation Signals.** In expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and/or any such sequence may be employed. Preferred embodiments include the bovine or human growth hormone polyadenylation signal, convenient and/or known to function well in various target cells. Also contemplated as an element of the expression cassette is a transcriptional termination site. These elements can serve to enhance message levels and/or to minimize read through from the cassette into other sequences.

[0057] **Origins of Replication.** In order to propagate a vector in a host cell, it may contain one or more origins of replication sites (often termed "ori"), which is a specific nucleic acid sequence at which replication is initiated. Alternatively an autonomously replicating sequence (ARS) can be employed if the host cell is yeast.

[0058] Selectable and Screenable Markers. In certain embodiments of the invention, the cells that contain the nucleic acid construct of the present invention may be identified *in vitro* or *in vivo* by including a marker in the expression vector. Such markers would confer an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker, such as the antibiotic resistance gene on the plasmid constructs (such as kanamycin, ampicillin, gentamycin, tetracycline, or chloramphenicol).

[0059] Usually the inclusion of a drug selection marker aids in the cloning and identification of transformants, for example, genes that confer resistance to neomycin, puromycin, hygromycin, DHFR, GPT, zeocin and histidinol are useful selectable markers. In addition to markers conferring a phenotype that allows for the discrimination of transformants based on the implementation of conditions, other types of markers including screenable markers such as GFP, whose basis is colorimetric analysis, are also contemplated. Alternatively, screenable enzymes may be utilized. One of skill in the art would also know how to employ immunologic markers, possibly in conjunction with FACS analysis. The marker used is not believed to be important, so long as it is capable of being expressed simultaneously with the nucleic acid encoding a gene product. Further examples of selectable and screenable markers are well known to one of skill in the art.

[0060] Growth hormone releasing hormone: Growth hormone releasing hormone ("GHRH") has a short half-life in the circulatory system, both in humans and pigs (Frohman et al., 1984). However, by employing GHRH analogs with a prolonged biological half-life and/or improved secretagogue activity, it was possible to achieve enhanced growth hormone ("GH") secretion. Therefore, GHRH mutants were generated by site-directed mutagenesis of a porcine (1-40)OH form of the cDNA (Seq ID #1). The site directed mutagenesis altered several amino acid codons of the wild type porcine GHRH (Seq ID #2). TI-GHRH mutant composition is shown in Figure 1 and (Seq ID #3). The substitution of Gly15 to Ala15 was used to increase α -helical conformation and

amphiphilic structure that resulted in decreased cleavage by trypsin-like enzymes (Su et al., 1991). Also, GHRH analogs with the Ala15-substitution display a 4-5 times higher affinity for the GHRH receptor (Reiss et al., 1993). We substituted Met27, Ser28 with Leu27, Asn28 (Kubiak et al., 1989) in order to reduce loss of biological activity due to oxidation of the Met27, thus forming a triple amino acid substitution denoted as 15/27/28-GHRH (Seq ID #4). Dipeptidyl peptidase IV is the prime serum GHRH degradative enzyme (Martin et al., 1993). Lower affinity dipeptidase substrates were created by further substitutions of 15/27/28-GHRH, and converting Ala2 for Ile2 (TI-GHRH, T1I2A15L27N28) or for Val2 (TV-GHRH – Seq ID #5) or by converting Tyr1 and Ala2 with His1 and Val2 (HV-GHRH – Seq ID #6). The HV-GHRH super-analog was presented in the U.S. Patent Application S.N. 10/021,403 filed on December 12, 2001 and titled “Administration of nucleic acid sequence to female animal to enhance growth in offspring” with Schwartz, et al., listed as inventors, and U.S. Patent Application S.N. 09/624,268 filed on July 24, 2000, 2000 and titled “Super Active Porcine Growth Hormone Releasing Hormone Analog” with Schwartz, et al., listed as inventors.

[0061] The invention may be better understood with reference to the following examples, which are representative of some of the embodiments of the invention, and are not intended to limit the invention.

EXAMPLE 1

[0062] To test the biological potency of the mutated porcine GHRH (pGHRH) cDNA, a plasmid vector was engineered that could direct very high levels of skeletal muscle-specific gene expression with the use of a newly described synthetic muscle promoter, SPc5-12 (Li et al., 1999). A 228 -bp fragment of pGHRH, which encodes the 31 amino acid signal peptide and a mature peptide pGHRH (Tyr1-Gly40) and/or the modified GHRH, or functional biological equivalents thereof, followed by the 3' untranslated region of the human GH (“hGH”) cDNA. All expression vector elements were operatively linked and incorporated into myogenic GHRH expression vectors. Skeletal myoblasts were transfected with each construct. Purified GHRH moieties from conditioned culture media were assayed for potency by their ability to induce GH secretion in pig anterior pituitary cell cultures. As shown in Figure 2, media were collected from the pituitary cell cultures after 24 hours and analyzed for porcine-specific GH by

radioimmunoassay. The modified GHRH species (15/27/28-GHRH; TI-GHRH; TV-GHRH, HV-GHRH) showed 20% to 50% improvements in their capacity to stimulate GH secretion compared to wild-type porcine GHRH ("wt-GHRH"), as indicated by the increase in porcine GH levels from a baseline value of 200ng/ml to 1600ng/ml. Although not wanting to be bound by theory, the increase was probably produced by an increased affinity for the GHRH receptors present on the pituitary cells.

[0063] Experimental Cell culture conditions were as follows: The Minimal Essential Medium ("MEM"), heat-inactivated horse serum ("HIHS"), gentamycin, Hanks Balanced Salt Solution ("HBSS"), lipofectamine were obtained from Gibco BRL (Grand Island, NY). Primary chicken myoblast cultures were obtained and transfected as previously described (Bergsma et al., 1986; Draghia-Akli et al., 1997). After transfection, the medium was changed to MEM which contained 2% HIHS to allow the cells to differentiate. Media and cells were harvested 72 hours post-differentiation. One day before harvesting, cells were washed twice in HBSS and the media changed to MEM, 0.1% bovine serum albumin ("BSA"). Conditioned media was treated by adding 0.25 volume of 1% trifluoroacetic acid ("TFA") and 1mM phenylmethylsulfonylflouride ("PMSF"), frozen at -80°C, lyophilized, purified on C-18 Sep-Columns (Peninsula Laboratories, Belmont, CA), relyophilized and used in the radioimmuno assay ("RIA") or resuspended in media conditioned for primary pig anterior pituitary culture. The pig anterior pituitary culture was obtained as previously described (Tanner et al., 1990).

[0064] Plasmid vectors containing the muscle specific synthetic promoter SPc5-12 were previously described (Li et al., 1999). Wild type and mutated porcine GHRH cDNAs were generated by site directed mutagenesis of GHRH cDNA (Altered Sites II *in vitro* Mutagenesis System, Promega, Madison, WI), and cloned into the BamHI/Hind III sites of pSPc5-12, to generate pSP-wt GHRH, or pSP-TI-GHRH respectively. The 3' untranslated region (3'UTR) of growth hormone was cloned downstream of GHRH cDNA. The resultant plasmids contained mutated coding region for GHRH, and the resultant amino acid sequences were not naturally present in mammals. Although not wanting to be bound by theory, the effects on treating GH deficient diseases is determined ultimately by the circulating levels of needed hormones. Several different plasmids that encode modified GHRH or functional biological equivalent thereof are as follows:

<u>Plasmid</u>	<u>Encoded Amino Acid Sequence</u>
wt-GHRH	YADAIFTNSYRKVLGQLSARKLLQDIMSRQQGERNQEQGA-OH
HV-GHRH	HVDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH
TI-GHRH	YIDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH
TV-GHRH	YVDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH
15/27/28GHRI	YADAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQGA-OH

In general, the encoded GHRH or functional biological equivalent thereof is of formula:



wherein: A₁ is a D-or L-isomer of an amino acid selected from the group consisting of tyrosine ("Y"), or histidine ("H"); A₂ is a D-or L-isomer of an amino acid selected from the group consisting of alanine ("A"), valine ("V"), or isoleucine ("I"); A₃ is a D-or L-isomer of an amino acid selected from the group consisting of alanine ("A") or glycine ("G"); A₄ is a D-or L-isomer of an amino acid selected from the group consisting of methionine ("M"), or leucine ("L"); A₅ is a D-or L-isomer of an amino acid selected from the group consisting of serine ("S") or asparagine ("N").

[0065] Another plasmid that was utilized included the pSP-SEAP construct that contains the SacI/ HindIII SPc5-12 fragment, secreted embryonic alkaline phosphatase (SEAP) gene and SV40 3'UTR from pSEAP-2 Basic Vector (Clontech Laboratories, Inc., Palo Alto, CA).

[0066] The plasmids described above do not contain polylinker, IGF-I gene, a skeletal alpha-actin promoter or a skeletal alpha actin 3' UTR /NCR. Furthermore, these plasmids were introduced by muscle injection, followed by *in vivo* electroporation, as described below.

[0067] In terms of "functional biological equivalents," it is well understood by the skilled artisan that, inherent in the definition of a "biologically functional equivalent" protein and/or polynucleotide, is the concept that there is a limit to the number of changes that may be made within a defined portion of the molecule while retaining a molecule with an acceptable level of equivalent biological activity. Functional biological equivalents are thus defined herein as those proteins (and polynucleotides) in selected amino acids (or codons) may be substituted. A peptide comprising a functional biological equivalent of GHRH is a polypeptide that has been engineered to contain distinct amino

acid sequences while simultaneously having similar or improved biological activity when compared to GHRH. For example one biological activity of GHRH is to facilitate growth hormone ("GH") secretion in the subject.

EXAMPLE 2

[0068] Stability of wild type GHRH and the analog TI-GHRH was then tested in porcine plasma by incubation of GHRH peptides in plasma, followed by solid phase extraction and HPLC analysis (Su et al., 1991). As shown in Figure 3, at least 60% of the pGHRH was degraded within 30 minutes of incubation in plasma. In contrast, incubation of TI-GHRH in pig plasma for up to 6 hours showed that at least 70% of the polypeptide was protected against enzymatic cleavage indicating a considerable increase in the resistance of TI-GHRH to serum protease activity. No degradation was seen in the first 30 minutes of incubation. *Plasma proteolytic activity on GHRH molecules.* Chemically synthesized TI-GHRH was prepared by peptide synthesis. Briefly, pooled porcine plasma was collected from control pigs, and stored at -80°C. At the time of the test, the porcine plasma was thawed, centrifuged and allowed to equilibrate at 37°C. Mutant and wild-type GHRH samples were dissolved in the plasma sample to a final concentration of 300µg/ml. Immediately after the addition of the GHRH, and 15, 30, 120 and 240 minutes later, 1mL of plasma was withdrawn and acidified with 1mL of 1M TFA. Acidified plasma was purified on C-18 affinity SEP-Pak columns, lyophilized, and analyzed by HPLC, using a Waters 600 multi-system delivery system, a Waters intelligent sample processor, type 717 and a Waters spectromonitor 490 (Waters Associates, Millipore Corp., Milford, MA). The mobile phase was (A) 0.1%TFA in H₂O, (B) 0.1%TFA in 95% ACN and 5% H₂O; the gradient was 80% (B) in 30 minutes. The flow rate was 0.75mL/min. Detection was performed at 214nm. The percent of peptide degraded at these time points was measured by integrated peak measurements.

EXAMPLE 3

[0069] **Biological activity of TI-GHRH in young animals.** Muscle injection of pSP-TI-GHRH increases IGF-I serum levels over two months in treated mice. We asked if the optimized protease resistant pSP-TI-GHRH vector could affect *in vivo*, long-term expression of GHRH and stimulate secretion of GH and subsequently, IGF-I. Schematic maps of pSP-TI-GHRH, the wild type construct, pSP-wt GHRH (positive

control), and an *E.coli*. β -galactosidase expression vector, pSP- β al (placebo control), are shown in Figure 4. Five weeks old SCID mice (immuno-deficient mice) were injected into the tibialis anterior muscle with 7.5 micrograms of one of the constructs. The injected muscle was placed within a caliper and electroporated, using optimized conditions of 200V/cm with 6 pulses of 60 milliseconds, as described in Materials and Methods (Aihara and Miyazaki, 1998).

[0070] Animals were bled up to two month post-injection and serum was used for IGF-I measurements (Figure 5). At 14 and 28 days post-injection, blood was collected and IGF-I levels were measured. All GHRH plasmid injected groups had highly significant increases in IGF-I levels compared to control animals, $p < 0.005$. Some groups developed neutralizing antibodies, and in these cases the IGF-I levels dropped by the second time point. The animals injected with TI-GHRH did not develop any antibodies, and their modified TI-GHRH expression continued for two month, and correlated with significant changes in their body composition.

[0071] At the end of the experiment, body composition was performed *in vivo*, using a dual x-ray absorptiometry technique (DEXA), with a high resolution scanner - PIXImus, and than post-mortem at necropsy. Blood was collected, centrifuged immediately at 4°C, and stored at -80°C prior to analysis. Organs, carcass, fat from injected animals and controls were removed, weighed and snap frozen in liquid nitrogen. The TI-GHRH ($p < 0.03$) and HV-GHRH injected animals were significantly heavier than controls at the end of the experiment (Figure 6).

[0072] Body composition analysis by DEXA (total body fat, non-bone lean tissue mass and bone mineral area, content and density) showed significant changes in animals injected with the TI-GHRH plasmid. Non-bone lean body mass (Figure 7) increased by 11% in TI-GHRH treated animals versus controls ($p < 0.036$). The HV-GHRH injected animals had a significant increase in lean body mass of almost 5%, as seen in previous experiments.

[0073] Upon injection with the TI-GHRH encoding plasmid, significant changes occurred in bone mineral area (Figure 8), that increased by 10.7%, ($p < 0.027$).

Example 4

[0074] A long-lasting therapy has the potential to replace classical GH therapy regimens and may stimulate the GH axis in a more physiologically appropriate manner. It is known that GHRH stimulates bone formation (Dubreuil et al., 1996), and our therapy may be used to promote post-fracture bone growth. Data show that GH plus IGF-I (delivered as recombinant proteins) synergistically increase lean muscle and body weight, total body weight, and were more effective in re-epithelialization of a burn wound than either GH or IGF-I alone (Meyer et al., 1996). Studies also showed that long-term stimulation of the GH axis, which includes doses in the range given to humans during clinical trials of GH deficiency and to revert age-related physiologic declines, has no overt deleterious effects on longevity and pathology in aged rodents (Kalu et al., 1998).

[0075] Using modified GHRH or functional biological equivalents thereof, it was possible to show that body composition was altered in treated animals. For example, twenty nine ("29") month old mice were injected into the tibialis anterior muscle with 15 micrograms TI-GHRH plasmid. The control group received the pSP- β gal construct previously described. Injection followed by electroporation with calipers with standard conditions 200V/cm, gap 5 mm on average, 3 pulses x 2, 50 milliseconds/pulse. Cardiac function in these animals was evaluated by Doppler at day 0, 10 and 20 and echocardiogram at day 20. At the end of the experiment, body composition was performed post-mortem.

[0076] Body length had a tendency to be increased in treated animals, suggestive of axial growth ($p < 0.07$) (Figure 9). Furthermore, tibia length / total body weight was significantly increased in TI-GHRH animals ($p < 0.006$) (Figure 10), a sign of bone remodeling and growth in these extremely old animals.

[0077] Aging, treated animals had reduced spleen weights / total body weight at necropsy (Figure 11). Although not wanting to be bound by theory, this observed reduction was probably due to lymphocyte mobilization from the lymphatic organ to the general circulation. This change is sign of increased immune surveillance. Growth hormone ("GH") is known to enhance immune responses, whether directly or through the IGF-I. GH secretagogues ("GHS") have important effects on immunological functions in young and old mammals. In young mice, GHS cause a significant increase in

peripheral blood lymphocyte counts. Old mice, treated with GHS for 3 weeks show significant resistance to the initiation of transplanted tumors and the subsequent metastases (Koo et al., 2001).

[0078] IGF-I also modulates the immune function, and has two major effects on B cell development: it acts as a differentiation factor to potentiate pro-B to pre-B cell maturation (Landreth et al., 1992), and it acts as a B cell proliferation cofactor to synergize with IL-7 (Landreth et al., 1985). There is evidence that macrophages are a rich source of IGF-I and that bone marrow stromal cells also produce IGF binding proteins ("IGFBP") (Abboud et al., 1991). The treatment of mice with recombinant IGF-I increased the number of pre-B and mature B cells in bone marrow (Jardieu et al., 1994). The mature B cell remains sensitive to IGF-I as immunoglobulin production is also stimulated by IGF-I *in vitro* and *in vivo* (Robbins et al., 1994). The administration of recombinant IGF-I has been shown to increase the size of lymphoid organs in other species. In 1-yr-old sheep, an 8-week regimen of three daily injections of recombinant IGF-I increased spleen weight by 40% (Cottam et al., 1992). In the rabbit, cat, and dog similar effects of IGF-I have been observed. In the rhesus monkey, IGF-I also expands lymphocyte numbers (LeRoith et al., 1996). In blood, the percent CD4 cell count and the CD4/CD8 ratio falls with IGF-I treatment but were normalized by GH plus IGF-I. In the spleen combination treatment almost triples the percent CD4 cells and more than doubles the CD4/CD8 ratio. This paradox of differential effects on lymphocyte populations in different body compartments may be due to the anabolic hormones affecting lymphocyte trafficking as GH and IGF-I appears to cause lymphocytes to accumulate in lymphoid organs at the expense of lymphocyte numbers in the circulation (Clark et al., 1993; Jardieu et al., 1994). These observations in primates make it more likely that IGF-I will prove useful in other species such as dogs, cats or humans to improve immune function, especially after damage to the immune system or in immune senescence in the elderly (Auernhammer and Strasburger, 1995) or in patients with cancer.

[0079] TI-GHRH treated animals also had increased cardiac function, as assessed by Doppler and echocardiogram. Heart rate is stable (within 5% throughout the assay) (Figure 12A). Peak aortic flow velocity does not change either (Figure 12B). This is unexpected as GH and IGF-I are supposed to induce eNOS and reduce peripheral vascular

resistance. Altered diastolic filling is an important contributor to several cardiovascular disorders (Houlind et al., 2002; Richartz et al., 2002; Tang et al., 2002). Peak early filling velocity, a measure of the cardiac diastolic filling indices, increases impressively by 20% at 10 days post-treatment (Figure 12C).

[0080] Among the non-viral techniques for plasmid transfer *in vivo*, the direct injection of plasmid DNA into muscle is simple, inexpensive, and safe. Applications of this methodology have been limited by the relatively low expression levels of the transferred DNA expression vectors. Previously, these levels have been insufficient to ensure systemic physiological levels of secreted proteins such as hormones, neurotrophic factors or coagulation factors in large mammals. Although not wanting to be bound by theory, in order to obtain growth of a large mammal by plasmid therapy, it is necessary to increase the potency of the myogenic vector system. The inventors recently described (Li et al., 1999) a strategy for the construction and the characterization of novel muscle synthetic promoters by the random assembly of E-boxes, MEF-2, TEF-1 and SRE sites. Several synthetic muscle promoters were identified whose transcriptional activity in terminally differentiated muscle greatly exceeded that of the natural myogenic skeletal α -actin gene promoter and viral promoters. Analysis of direct intramuscular injection of SPc5-12 driven DNA plasmid in normal mouse muscle revealed a 6-8-fold increase in activity over the ubiquitously expressed CMV promoter even after a month. As shown in figure 2, SPc5-12 was capable of eliciting moderate increases in growth and IGF-I levels by driving GHRH production in animals. Severe combined immunodeficient (SCID) adult male mice (aged 5-6 weeks at the beginning of the experiment) or NIH C57/Bl6 mice (aged 29 month) were housed and cared for in the animal facility of Baylor College of Medicine, Houston, TX. Animals were maintained under environmental conditions of 10h light / 14h darkness, in accordance with NIH Guidelines, USDA and Animal Welfare Act guidelines, and the protocol was approved by the Institutional Animal Care and Use Committee. On day 0, the animals were weighed and then, the left tibialis anterior muscle of mice was injected with plasmids in 25 μ l PBS. The injection was followed by caliper electroporation, as previously described (Draghia-Akli et al., 1999). The animals were bled periodically, and serum was used to measure IGF-I levels. At the end of the experiment, body composition was performed *in vivo*, using the DEXA technique, and than at necropsy. Blood was collected, centrifuged immediately at 4°C, and stored at -

80°C prior to analysis. Organs, carcass, and fat from injected animals and controls were removed, weighed and snap frozen in liquid nitrogen.

[0081] Mouse IGF-I Radioimmunoassay: Mouse IGF-I was measured by heterologous, 100% cross-reacting rat radioimmunoassay. The sensitivity of the assay was 0.8 ng/ml; intra-assay and inter-assay variation was 3.4% and 4.5% respectively. The statistics and values shown in the figures are the mean \pm s.e.m. Specific p values were obtained by comparison using Students t-test or ANOVA analysis. A $p < 0.05$ was set as the level of statistical significance.

[0082] Another significant improvement of our plasmid vector was the employment of a novel GHRH analog, TI-GHRH. Some individual amino acid substitutions leading to protease resistant GHRH molecules were previously tested in farm animals and humans (Frohman et al., 1989; Martin et al., 1993). The inventors have found that novel combination of five amino acid substitutions in the TI-GHRH construct resulted in increased GH secretagogue activity (as shown in assays on pig anterior pituitary somatotrophic cells) and was more resistant to serum proteases *in vivo* (Figure 2).

EXAMPLE 5

[0083] Although not wanting to be bound by theory, electro-plasmid therapy allows genes to be efficiently transferred and expressed in desired organs or tissues, and it may represent a new approach for highly effective plasmid supplementation therapy, that does not require viral genes or particles. The electroporation system was used previously in rodents and small animals and does not appear to cause significant distress. Electro-plasmid therapy increased transfection efficiency over 100-fold compared to classical plasmid therapy techniques and allowed for prolonged TI-GHRH expression.

[0084] Although not wanting to be bound by theory, enhanced biological potency and enhanced delivery protocols reduces the theoretical quantity of GHRH plasmid needed to achieve physiological levels of GH production and secretion. Treated mice did not experience any side effects from the therapy, had normal biochemical profiles, and with no associated pathology. The profound increases in IGF-I levels growth enhancement, the changes in body composition, increases in immune function and improved cardiac function indicate that ectopic expression of myogenic TI-GHRH vectors

has the potential to replace classical GH therapy regimens and may stimulate the GH axis in a more physiologically appropriate manner. The TI-GHRH molecule, which displays a high degree of stability and GH secretory activity, may also be useful in human clinical medicine, since the serum proteases that degrade GHRH are similar in most mammals.

[0085] Hormones (e.g. GHRH and GH) often contain a complex feedback-regulated pathway, which are further complicated by chronic conditions such as cancer or AIDS. Without direct experimentation of GHRH or biological equivalents used in plasmid mediated supplementation, a beneficial therapy could not have been predicted by one skilled in the art to determine which modified GHRH or functional biological equivalent thereof, encoded sequences will yield desired results. The invention described herein contains the compositions, descriptions, and results of essential experimentation that explored tissue specific and inducible regulation of distinctive nucleic acid sequences that encoded modified GHRH or biological equivalent thereof, which was not obvious based upon prior art.

[0086] One skilled in the art readily appreciates that the disclosed invention is well adapted to carry out the mentioned and inherent objectives. Growth hormone, growth hormone releasing hormone, modified growth hormone releasing hormone or functional biological equivalents, plasmids, vectors, pharmaceutical compositions, treatments, methods, procedures and techniques described herein are presented as representative of the preferred embodiments and are not intended as limitations of the scope of the invention. Thus, other uses will occur to those skilled in the art that are encompassed within the spirit and scope of the described invention.

The following documents and publications are incorporated by reference herein.

REFERENCES CITED

U.S. PATENT DOCUMENTS

- U.S. Patent No. 5,847,066 issued on December 8, 1998 with Coy et al. listed as inventors.
- U.S. Patent No. 5,846,936 issued on December 8, 1998 with Felix et al. listed as inventors.
- U.S. Patent No. 5,792,747 issued on August 11, 1998 with Schally et al. listed as inventors.
- U.S. Patent No. 5,776,901 issued on July 7, 1998 with Bowers et al. listed as inventors.
- U.S. Patent No. 5,756,264 issued on May 26, 1998 with Schwartz et al. listed as inventors.
- U.S. Patent No. 5,696,089 issued on December 9, 1997 with Felix et al. listed as inventors.
- U.S. Patent No. 5,486,505 issued on January 23, 1996 with Bowers et al. listed as inventors.
- U.S. Patent No. 5,137,872 issued on August 11, 1992 with Seely et al. listed as inventors.
- U.S. Patent No. 5,084,442 issued on January 28, 1992 with Felix et al. listed as inventors.
- U.S. Patent No. 5,036,045 issued on July 30, 1991 with Thorner listed as the inventor.
- U.S. Patent No. 5,023,322 issued on June 11, 1991 with Kovacs et al. listed as inventors.
- U.S. Patent No. 4,839,344 issued on June 13, 1989 with Bowers et al. listed as inventors.
- U.S. Patent No. 4,410,512 issued on October 18, 1983 with Bowers et al. listed as inventors.
- U.S. Patent No. RE33,699 issued on September 24, 1991 with Drenkler listed as the inventor.
- U.S. Patent No. 4,833,166 issued on May 23, 1989 with Grosvenor et al. listed as inventors.
- U.S. Patent No. 4,228,158 issued on October 14, 1980 with Momany et al. listed as inventors.
- U.S. Patent No. 4,228,156 issued on October 14, 1980 with Momany et al. listed as inventors.
- U.S. Patent No. 4,226,857 issued on October 7, 1980 with Momany et al. listed as inventors.

U.S. Patent No. 4,224,316 issued on September 23, 1980 with Momany et al. listed as inventors.

U.S. Patent No. 4,223,021 issued on September 16, 1980 with Momany et al. listed as inventors.

U.S. Patent No. 4,223,020 issued on September 16, 1980 with Momany et al. listed as inventors.

U.S. Patent No. 4,223,019 issued on September 16, 1980 with Momany et al. listed as inventors.

PUBLICATION REFERENCE LIST

Abboud, S. L., C. R. Bethel, and D. C. Aron. 1991. Secretion of insulin like growth factor I and insulin-like growth factor-binding proteins by murine bone marrow stromal cells. *J Clin. Invest* 88:470-475.

Acsadi, G., G. Dickson, D. R. Love, A. Jani, F. S. Walsh, A. Gurusinghe, Wolff, JA, and K. E. Davies. 1991. Human dystrophin expression in mdx mice after intramuscular injection of DNA constructs. *Nature* 352:815-818.

Aihara, H. and J. Miyazaki. 1998. Gene transfer into muscle by electroporation in vivo. *Nat. Biotechnol.* 16:867-870.

Auernhammer, C. J. and C. J. Strasburger. 1995. Effects of growth hormone and insulin-like growth factor I on the immune system. *European Journal of Endocrinology* 133:635-645.

Bergsma, D. J., J. M. Grichnik, L. M. Gossett, and R. J. Schwartz. 1986. Delimitation and characterization of cis-acting DNA sequences required for the regulated expression and transcriptional control of the chicken skeletal alpha-actin gene. *Molecular & Cellular Biology* 6:2462-2475.

Butler, A. A., G. R. Ambler, B. H. Breier, D. LeRoith, C. T. Roberts, Jr., and P. D. Gluckman. 1994. Growth hormone (GH) and insulin-like growth factor-I (IGF-I) treatment of the GH-deficient dwarf rat: differential effects on IGF-I transcription start site expression in hepatic and extrahepatic tissues and lack of effect on type I IGF receptor mRNA expression. *Mol. Cell Endocrinol.* 101:321-330.

Caroni, P. and C. Schneider. 1994. Signaling by insulin-like growth factors in paralyzed skeletal muscle: rapid induction of IGF1 expression in muscle fibers and prevention of interstitial cell proliferation by IGF-BP5 and IGF-BP4. *J. Neurosci.* 14:3378-3388.

Clark, R., J. Strasser, S. McCabe, K. Robbins, and P. Jardieu. 1993. Insulin-like growth factor-1 stimulation of lymphopoiesis. *J Clin. Invest* 92:540-548.

Corpas, E., S. M. Harman, M. A. Pineyro, R. Roberson, and M. R. Blackman. 1993. Continuous subcutaneous infusions of growth hormone (GH) releasing hormone 1-

3188856v13 108328/00067

- 44 for 14 days increase GH and insulin-like growth factor-I levels in old men. *Journal of Clinical Endocrinology & Metabolism* 76:134-138.
- Cottam, Y. H., H. T. Blair, B. W. Gallaher, R. W. Purchas, B. H. Breier, S. N. McCutcheon, and P. D. Gluckman. 1992. Body growth, carcass composition, and endocrine changes in lambs chronically treated with recombinantly derived insulin-like growth factor-I. *Endocrinology* 130:2924-2930.
- Danko, I. and J. A. Wolff. 1994. Direct gene transfer into muscle. *Vaccine* 12:1499-1502.
- Davis, H. L., R. G. Whalen, and B. A. Demeneix. 1993. Direct gene transfer into skeletal muscle in vivo: factors affecting efficiency of transfer and stability of expression. *Human Gene Therapy* 4:151-159.
- Draghia-Akli, R., M. L. Fiorotto, L. A. Hill, P. B. Malone, D. R. Deaver, and R. J. Schwartz. 1999. Myogenic expression of an injectable protease-resistant growth hormone-releasing hormone augments long-term growth in pigs. *Nat. Biotechnol.* 17:1179-1183.
- Draghia-Akli, R., X. G. Li, R. J. Schwartz. 1997. Enhanced Growth By Ectopic Expression Of Growth Hormone Releasing Hormone Using An Injectable Myogenic Vector. *nature biotechnology* 15:1285-1289.
- Draghia-Akli, R., P. B. Malone, L. A. Hill, K. M. Ellis, R. J. Schwartz, and J. L. Nordstrom. 2002. Enhanced animal growth via ligand-regulated GHRH myogenic-injectable vectors. *FASEB J.* 16:426-428.
- Dubreuil, P., T. Abribat, B. Broxup, and P. Brazeau. 1996. Long-term growth hormone-releasing factor administration on growth hormone, insulin-like growth factor-I concentrations, and bone healing in the Beagle. *Canadian Journal of Veterinary Research* 60:7-13.
- Etherton, T. D., J. P. Wiggins, C. S. Chung, C. M. Evock, J. F. Rebhun, and P. E. Walton. 1986. Stimulation of pig growth performance by porcine growth hormone and growth hormone-releasing factor. *Journal of Animal Science* 63:1389-1399.
- Foncea, R., M. Andersson, A. Ketterman, V. Blakesley, M. Sapag-Hagar, P. H. Sugden, D. LeRoith, and S. Lavandero. 1997. Insulin-like growth factor-I rapidly activates multiple signal transduction pathways in cultured rat cardiac myocytes. *J. Biol. Chem.* 272:19115-19124.
- Frohman, L. A., T. R. Downs, and P. Chomczynski. 1992. Regulation of growth hormone secretion. [Review]. *Frontiers in Neuroendocrinology* 13:344-405.
- Frohman, L. A., T. R. Downs, E. P. Heimer, and A. M. Felix. 1989. Dipeptidylpeptidase IV and trypsin-like enzymatic degradation of human growth hormone-releasing hormone in plasma. *J. Clin. Invest.* 83:1533-1540.
- Frohman, L. A., J. L. Thominet, C. B. Webb, M. L. Vance, H. Uderman, J. Rivier, W. Vale, and M. O. Thorner. 1984. Metabolic clearance and plasma disappearance
- 3188856v13 108328/00067

- rates of human pancreatic tumor growth hormone releasing factor in man. *J. Clin. Invest.* 73:1304-1311.
- Gesundheit, N. and J. K. Alexander. 1995. Endocrine Therapy with Recombinant Hormones and Growth Factors. In: B. D. Weintraub (Ed.) *Molecular Endocrinology: Basic Concepts and Clinical Correlations*. pp. 491-507. Raven Press, Ltd., New York.
- Hoess, R. H. and K. Abremski. 1985. Mechanism of strand cleavage and exchange in the Cre-lox site-specific recombination system. *J. Mol. Biol.* 181:351-362.
- Houlind, K., A. P. Schroeder, H. Stodkilde-Jorgensen, P. K. Paulsen, H. Egeblad, and E. M. Pedersen. 2002. Intraventricular dispersion and temporal delay of early left ventricular filling after acute myocardial infarction. Assessment by magnetic resonance velocity mapping. *Magn Reson. Imaging* 20:249-260.
- Jardieu, P., R. Clark, D. Mortensen, and K. Dorshkind. 1994. In vivo administration of insulin-like growth factor-I stimulates primary B lymphopoiesis and enhances lymphocyte recovery after bone marrow transplantation. *J. Immunol.* 152:4320-4327.
- Kalu, D. N., P. B. Orhii, C. Chen, D. Y. Lee, G. B. Hubbard, S. Lee, and Y. Olatunji-Bello. 1998. Aged-rodent models of long-term growth hormone therapy: lack of deleterious effect on longevity. *J. Gerontol. A. Biol. Sci. Med. Sci.* 53:B452-B463.
- Koo, G. C., C. Huang, R. Camacho, C. Trainor, J. T. Blake, A. Sirotina-Meisher, K. D. Schleim, T. J. Wu, K. Cheng, R. Nargund, and G. McKissick. 2001. Immune enhancing effect of a growth hormone secretagogue. *J. Immunol.* 166:4195-4201.
- Kubiak, T. M., C. R. Kelly, and L. F. Krabill. 1989. In vitro metabolic degradation of a bovine growth hormone-releasing factor analog Leu27-bGRF(1-29)NH₂ in bovine and porcine plasma. Correlation with plasma dipeptidylpeptidase activity. *Drug Metabolism & Disposition* 17:393-397.
- Landreth, K. S., D. Engelhard, M. H. Beare, P. W. Kincade, N. Kapoor, and R. A. Good. 1985. Regulation of human B lymphopoiesis: effect of a urinary activity associated with cyclic neutropenia. *J. Immunol.* 134:2305-2309.
- Landreth, K. S., R. Narayanan, and K. Dorshkind. 1992. Insulin-like growth factor-I regulates pro-B cell differentiation. *Blood* 80:1207-1212.
- Lapierre, H., G. Pelletier, D. Petitclerc, P. Dubreuil, J. Morisset, P. Gaudreau, Y. Couture, and P. Brazeau. 1991. Effect of human growth hormone-releasing factor and(or) thyrotropin-releasing factor on growth, carcass composition, diet digestibility, nutrient balance, and plasma constituents in dairy calves. *Journal of Animal Science* 69:587-598.
- LeRoith, D., J. Yanowski, E. P. Kaldjian, E. S. Jaffe, T. LeRoith, K. Purdue, B. D. Cooper, R. Pyle, and W. Adler. 1996. The effects of growth hormone and insulin-

like growth factor I on the immune system of aged female monkeys. *Endocrinology* 137:1071-1079.

Li, X., E. M. Eastman, R. J. Schwartz, and R. Draghia-Akli. Synthetic muscle promoters: activities exceeding naturally occurring regulatory sequences. *nature biotechnology* 17[3], 241-245. 1999.

Ref Type: Generic

Liu, J. L. and D. LeRoith. 1999. Insulin-like growth factor I is essential for postnatal growth in response to growth hormone. *Endocrinology* 140:5178-5184.

Lowe, W. L., Jr., M. Adamo, H. Werner, C. T. Roberts, Jr., and D. LeRoith. 1989. Regulation by fasting of rat insulin-like growth factor I and its receptor. Effects on gene expression and binding. *J. Clin. Invest* 84:619-626.

Martin, R. A., D. L. Cleary, D. M. Guido, H. A. Zurcher-Neely, and T. M. Kubiak. 1993. Dipeptidyl peptidase IV (DPP-IV) from pig kidney cleaves analogs of bovine growth hormone-releasing factor (bGRF) modified at position 2 with Ser, Thr or Val. Extended DPP-IV substrate specificity? *Biochimica et Biophysica Acta* 1164:252-260.

Meyer, N. A., R. E. Barrow, and D. N. Herndon. 1996. Combined insulin-like growth factor-1 and growth hormone improves weight loss and wound healing in burned rats. *J. Trauma* 41:1008-1012.

Muramatsu, T., A. Nakamura, and H. M. Park. 1998. In vivo electroporation: A powerful and convenient means of nonviral gene transfer to tissues of living animals (Review) [In Process Citation]. *Int. J. Mol. Med.* 1:55-62.

Parks, J. S., R. W. Pfaffle, M. R. Brown, H. Abdul-Latif, and L. R. Meacham. 1995. Growth Hormone Deficiency. In: B. D. Weintraub (Ed.) *Molecular Endocrinology: Basic Concepts and Clinical Correlations*. pp. 473-490. Raven Press, Ltd., New York.

Parrizas, M. and D. LeRoith. 1997. Insulin-like growth factor-1 inhibition of apoptosis is associated with increased expression of the bcl-xL gene product. *Endocrinology* 138:1355-1358.

Rabinovsky, E. D., G. M. Smith, D. P. Browder, H. D. Shine, and J. L. McManaman. 1992. Peripheral nerve injury down-regulates CNTF expression in adult rat sciatic nerves. *J. Neurosci. Res.* 31:188-192.

Reiss, K., J. Kajstura, J. M. Capasso, T. A. Marino, and P. Anversa. 1993. Impairment of myocyte contractility following coronary artery narrowing is associated with activation of the myocyte IGF1 autocrine system, enhanced expression of late growth related genes, DNA synthesis, and myocyte nuclear mitotic division in rats. *Exp. Cell Res.* 207:348-360.

- Richartz, B., G. Werner, M. Ferrari, and H. Figulla. 2002. Comparison of left ventricular systolic and diastolic function in patients with idiopathic dilated cardiomyopathy and mild heart failure versus those with severe heart failure. *Am. J. Cardiol.* 90:390.
- Robbins, K., S. McCabe, T. Scheiner, J. Strasser, R. Clark, and P. Jardieu. 1994. Immunological effects of insulin-like growth factor-I--enhancement of immunoglobulin synthesis. *Clin. Exp. Immunol.* 95:337-342.
- Su, C. M., L. R. Jensen, E. P. Heimer, A. M. Felix, Y. C. Pan, and T. F. Mowles. 1991. In vitro stability of growth hormone releasing factor (GRF) analogs in porcine plasma. *Hormone & Metabolic Research* 23:15-21.
- Tang, W., D. K. Arnett, R. B. Devereux, L. D. Atwood, D. W. Kitzman, and D. C. Rao. 2002. Linkage of left ventricular early diastolic peak filling velocity to chromosome 5 in hypertensive African Americans: the HyperGEN echocardiography study. *Am. J. Hypertens.* 15:621-627.
- Tanner, J. W., S. K. Davis, N. H. McArthur, J. T. French, and T. H. Welsh, Jr. 1990. Modulation of growth hormone (GH) secretion and GH mRNA levels by GH-releasing factor, somatostatin and secretagogues in cultured bovine adenohypophysial cells. *J. Endocrinol.* 125:109-115.
- Thorner, M. O., L. A. Frohman, D. A. Leong, J. Thominet, T. Downs, P. Hellmann, J. Chitwood, J. M. Vaughan, and W. Vale. 1984. Extrahypothalamic growth-hormone-releasing factor (GRF) secretion is a rare cause of acromegaly: plasma GRF levels in 177 acromegalic patients. *Journal of Clinical Endocrinology & Metabolism* 59:846-849.
- Tripathy, S. K., E. C. Svensson, H. B. Black, E. Goldwasser, M. Margalith, Hobart, PM, and J. M. Leiden. 1996. Long-term expression of erythropoietin in the systemic circulation of mice after intramuscular injection of a plasmid DNA vector. *Proc. Natl. Acad. Sci. USA* 93:10876-10880.
- Tsurumi, Y., S. Takeshita, D. Chen, M. Kearney, S. T. Rossow, J. Passeri, J. R. Horowitz, J. F. Symes, and J. M. Isner. 1996. Direct intramuscular gene transfer of naked DNA encoding vascular endothelial growth factor augments collateral development and tissue perfusion [see comments]. *Circulation* 94:3281-3290.
- van Rooij, E. M., B. L. Haagmans, H. L. Glansbeek, Y. E. de Visser, M. G. de Bruin, W. Boersma, and A. T. Bianchi. 2000. A DNA vaccine coding for glycoprotein B of pseudorabies virus induces cell-mediated immunity in pigs and reduces virus excretion early after infection. *Vet. Immunol. Immunopathol.* 74:121-136.
- Veldhuis, J. D., A. Iranmanesh, and A. Weltman. 1997. Elements In The Pathophysiology Of Diminished Growth Hormone (GH) Secretion In Aging Humans. *Endocrine* 7:41-48.

- Wang, L., W. Ma, R. Markovich, W. L. Lee, and P. H. Wang. 1998. Insulin-like growth factor I modulates induction of apoptotic signaling in H9C2 cardiac muscle cells. *Endocrinology* 139:1354-1360.
- Wolff, J. A., R. W. Malone, P. Williams, W. Chong, G. Acsadi, A. Jani, Felgner, and PL. 1990. Direct gene transfer into mouse muscle in vivo. *Science* 247:1465-1468.

WHAT IS CLAIMED IS:

1. A composition comprising: a modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof with a general formula:

~~-A₁-A₂-DAIFTNSYRKVL-A₃-QLSARKLLQDI-A₄-A₅-RQQGERNQEQGA-OH~~

wherein: A₁ is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H"); A₂ is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I"); A₃ is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G"); A₄ is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L"); A₅ is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N").

2. The composition of claim 1, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #3.

3. The composition of claim 1, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #4.

4. The composition of claim 1, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #5.

5. The composition of claim 1, wherein the modified GHRH is a biologically active polypeptide; and the functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to a wild-type GHRH ("wt-GHRH") polypeptide.

6. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof increases growth hormone ("GH") secretion when delivered into a subject.

7. The composition of claim 6, wherein the subject is a human or domesticated animal.

8. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof increases a insulin-like growth factor I ("IGF-I") secretion when delivered into a subject.
9. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof improves cardiac function when delivered into a subject.
10. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof increases weight gain when delivered into a subject.
11. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof increases bone mineral density when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
12. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof increases body length, when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
13. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof improves immune function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
14. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof improves cardiac function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
15. The composition of claim 5, wherein the modified GHRH or functional biological equivalent thereof is resistant to degradation when compared to the wt-GHRH.
16. The composition of claim 1, wherein the modified GHRH or functional biological equivalent thereof is translated from an encoded nucleic acid molecule, wherein the nucleic acid molecule is DNA or RNA.

17. The composition of claim 16, wherein the nucleic acid molecule further comprises a synthetic mammalian expression plasmid.

18. The composition of claim 17, wherein the synthetic mammalian expression plasmid further comprises:

- a synthetic or eukaryotic promoter;
- a poly-adenylation signal;
- a selectable marker gene promoter;
- a ribosomal binding site;
- a selectable marker gene sequence; and
- an origin of replication;

wherein the synthetic or eukaryotic promoter, the nucleic acid sequence encoding the modified GHRH or functional biological equivalent thereof, and the poly adenylation signal comprise therapeutic elements of the synthetic mammalian expression plasmid; the therapeutic elements are operatively linked and located in a first operatively-linked arrangement; the selectable marker gene promoter, the ribosomal binding site, the selectable marker gene sequence, and the origin of replication comprise replication elements of the synthetic mammalian expression plasmid; the replication elements are operatively linked and located in a second operatively-linked arrangement; the first-operatively-linked arrangement and the second-operatively-linked arrangement comprise a circular structure of the synthetic mammalian expression plasmid; and the synthetic mammalian expression plasmid is utilized for plasmid mediated gene supplementation.

19. The synthetic mammalian expression plasmid of claim 18, further comprising a 3' untranslated region ("UTR") operatively linked to the first operatively-linked arrangement. .

20. The synthetic mammalian expression plasmid of claim 19, wherein the 3' untranslated region ("UTR") comprises a portion of a human growth hormone 3'UTR.

21. The synthetic mammalian expression plasmid of claim 18, wherein the first operatively-linked arrangement comprises SeqID#7.
22. The synthetic mammalian expression plasmid of claim 18, wherein the first operatively-linked arrangement comprises SeqID#8.
23. The synthetic mammalian expression plasmid of claim 18, wherein the first operatively-linked arrangement comprises SeqID#9.
24. A nucleic acid expression construct encoding a modified growth hormone releasing hormone ("GHRH") or functional biological equivalent thereof with a general formula:
-A₁-A₂-DAIFTNSYRKVL-A₃-QLSARKLLQDI-A₄-A₅-RQQGERNQEQGA-OH
wherein: A₁ is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H"); A₂ is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I"); A₃ is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G"); A₄ is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L"); A₅ is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N"), and wherein the modified GHRH.
25. The composition of claim 24, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #3.
26. The composition of claim 24, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #4.
27. The composition of claim 24, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #5.
28. The composition of claim 24, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.
29. The composition of claim 28, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.

30. The composition of claim 28, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

31. The composition of claim 24, wherein the nucleic acid expression construct initiates expression of the encoded modified GHRH or functional biological equivalent thereof when delivered into a cell of a subject.

32. The composition of claim 31, wherein the encoded modified GHRH or functional biological equivalent thereof is capable of being expressed in a tissue specific cell of the subject.

33. The composition of claim 32, wherein the tissue specific cell of the subject comprises a muscle cell.

34. The composition of claim 31, wherein the subject is a human or domesticated animal.

35. The composition of claim 24, wherein the encoded modified GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biologically activity when compared to the GHRH polypeptide.

36. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof facilitates growth hormone ("GH") secretion when delivered into a subject.

37. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof increases a insulin-like growth factor I ("IGF-I") secretion when delivered into a subject.

38. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof increases weight gain when delivered into a subject.

39. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof increases bone mineral density when the encoded modified GHRH or functional biological equivalent thereof is delivered into a subject.
40. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof increases body length when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
41. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof improves immune function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
42. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof improves cardiac function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.
43. The composition of claim 24, wherein the encoded modified GHRH or functional biological equivalent thereof is resistant to degradation when compared to the wt GHRH.
44. A composition comprising: a modified GHRH of Seq ID #3.
45. The composition of claim 44, wherein the modified GHRH of Seq ID #3 is encoded by a nucleic acid expression construct.
46. The composition of claim 45, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.
47. The composition of claim 46, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.
48. The composition of claim 46, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.

49. A composition comprising: a modified GHRH of Seq ID #4.
50. The composition of claim 49, wherein the modified GHRH of Seq ID #4 is encoded by a nucleic acid expression construct.
51. The composition of claim 50, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.
52. The composition of claim 51, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.
53. The composition of claim 51, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.
54. A composition comprising: a modified GHRH of Seq ID #5.
55. The composition of claim 54, wherein the modified GHRH of Seq ID #5 is encoded by a nucleic acid expression construct.
56. The composition of claim 55, wherein the nucleic acid expression construct further comprises, a transfection-facilitating polypeptide.
57. The composition of claim 56, wherein the transfection-facilitating polypeptide comprises a charged polypeptide.
58. The composition of claim 56, wherein the transfection-facilitating polypeptide comprises poly-L-glutamate.
59. A method of increasing growth hormone ("GH") secretion in a subject comprising: delivering into a cell of the subject a nucleic acid expression construct that encodes a modified growth-hormone-releasing-hormone ("GHRH") or functional biological equivalent thereof, wherein the modified GHRH or functional biological equivalent thereof is of formula:

$$-A_1-A_2-DAIFTSYRKVL-A_3-QLSARKLLQDI-A_4-A_5-RQQGERNQEQGA-OH$$

3188856v13 108328/00067

wherein: A₁ is a D-or L-isomer of the amino acid tyrosine ("Y"), or histidine ("H"); A₂ is a D-or L-isomer of the amino acid alanine ("A"), valine ("V"), or isoleucine ("I"); A₃ is a D-or L-isomer of the amino acid alanine ("A") or glycine ("G"); A₄ is a D-or L-isomer of the amino acid methionine ("M"), or leucine ("L"); A₅ is a D-or L-isomer of the amino acid serine ("S") or asparagine ("N").

60. The method of claim 59, wherein delivering into the cell of the subject the nucleic acid expression construct is via electroporation.

61. The method of claim 59, wherein the cell of the subject is a somatic cell, a stem cell, or a germ cell.

62. The method of claim 59, wherein the cell of the subject is a muscle cell.

63. The method of claim 59, wherein the delivering into the cell of the subject the nucleic acid expression construct initiates expression of the encoded modified GHRH or functional biological equivalent thereof.

64. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is expressed in a tissue specific cell of the subject.

65. The method of claim 59, wherein the modified GHRH is a biologically active polypeptide; and the encoded functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to the GHRH polypeptide.

66. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #3.

67. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #4.

68. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is Seq ID #5.

69. The method of claim 59, wherein the modified GHRH is a biologically active polypeptide; and the functional biological equivalent of GHRH is a polypeptide that has been engineered to contain a distinct amino acid sequence while simultaneously having similar or improved biological activity when compared to a wild-type GHRH ("wt-GHRH") polypeptide.

70. The method of claim 59, wherein the subject is a human or domesticated animal.

71. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof increases insulin-like growth factor I ("IGF-I") secretion when delivered into a subject.

72. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof increases weight gain when delivered into a subject.

73. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof increases bone mineral density when the modified GHRH or functional biological equivalent thereof is delivered into a subject.

74. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof increases body length, when the modified GHRH or functional biological equivalent thereof is delivered into a subject.

75. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof improves the immune function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.

76. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof improves cardiac function when the modified GHRH or functional biological equivalent thereof is delivered into a subject.

77. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is resistant to degradation when compared to the wt-GHRH.

78. The method of claim 59, wherein the modified GHRH or functional biological equivalent thereof is resistant to degradation when compared to the wt-GHRH.

1/12

YADAIFTNSYRKVLGQLSARKLLQDIMSRQQGERNQEQA-OH porcine wild-type
YIDAIFTNSYRKVLAQLSARKLLQDILNRQQGERNQEQA-OH TI-GHRH

Figure 1




Figure 2

3/12




Figure 3





Figure 4



B.

Figure 5

6/12




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

12/12




Figure 12

Sequence Listing.ST25.txt
SEQUENCE LISTING

<110> Advisys
Baylor College of Medicine
Baylor

<120> PROTEASE RESISTANT TI-GROWTH HORMONE RELEASING HORMONE ("GHRH")

<130> 108328.00067 - AVSI-0024

<160> 9

<170> PatentIn version 3.1

<210> 1
<211> 219
<212> DNA
<213> artificial sequence

<220>
<223> This is the cDNA for Porcine growth hormone releasing hormone

<400> 1
atggtgctct ggggtgttctt ctttgtgatc ctcaccctca gcaacagctc ccactgctcc 60
ccacctcccc ctttgaccct caggatgcgg cggcacgtag atgccatctt caccaacagc 120
taccggaagg tgctggccca gctgtccgcc cgcaagctgc tccaggacat cctgaacagg 180
cagcagggag agaggaacca agagcaagga gcataatga 219

<210> 2
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is the amino acid sequence for porcine growth hormone releasing hormone.

<400> 2
Tyr Ala Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Gly Gln
1 5 10 15
Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Met Ser Arg Gln Gln Gly
20 25 30
Glu Arg Asn Gln Glu Gln Gly Ala
35 40

<210> 3
<211> 40
<212> PRT
<213> artificial sequence

<220>
<223> This is a modified amino acid sequence for growth hormone releasing hormone (GHRH). alpha-helical conformation was increased by s

Sequence Listing.ST25.txt
 ubstituting cly15 to Ala15.

<400> 3

Tyr Ile Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
 1 5 10 15
 Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
 20 25 30
 Glu Arg Asn Gln Glu Gln Gly Ala
 35 40

<210> 4

<211> 40

<212> PRT

<213> artificial sequence

<220>

<223> This is a porcine growth hormone releasing hormone ("GHRH") that
 has the following substitutions: Met27, Ser28 with Leu27 and Asn28.

<400> 4

Tyr Ala Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
 1 5 10 15
 Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
 20 25 30
 Glu Arg Asn Gln Glu Gln Gly Ala
 35 40

<210> 5

<211> 40

<212> PRT

<213> artificial sequence

<220>

<223> This is a growth hormone releasing hormone that has a val2 substitution for a ile 2.

<400> 5

Tyr Val Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
 1 5 10 15
 Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
 20 25 30
 Glu Arg Asn Gln Glu Gln Gly Ala
 35 40

Sequence Listing.ST25.txt

<210> 6
 <211> 40
 <212> PRT
 <213> artificial sequence

<220>
 <223> This is a growth hormone releasing hormone ("GHRH") with a His1 and val2 substituting the Try1 and Ala2.

<400> 6

His Val Asp Ala Ile Phe Thr Asn Ser Tyr Arg Lys Val Leu Ala Gln
 1 5 10 15

Leu Ser Ala Arg Lys Leu Leu Gln Asp Ile Leu Asn Arg Gln Gln Gly
 20 25 30

Glu Arg Asn Gln Glu Gln Gly Ala
 35 40

<210> 7
 <211> 3534
 <212> DNA
 <213> artificial sequence

<220>
 <223> This is the nucleic acid sequence for the operatively linked components of the TI-GHRH plasmid.

<400> 7

gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60

accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120

gtgaggaatg gtggggagtt atttttagag cggtgaggaa ggtgggcagg cagcagggtgt 180

tggcgctcta aaaataactc ccgggagtta tttttagagc ggaggaatgg tggacaccca 240

aatatggcga cggttcctca cccgtcgcca tatgtgggtg tccgccctcg gccggggccg 300

cattcctggg ggccgggagg tgctcccggc cgcctcgata aaagggtccg gggccggagg 360

cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420

ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480

ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tcccactgct cccacacctc 540

ccctttgacc ctcaggatgc ggcggtatat cgatgccatc ttcaccaaca gctaccggaa 600

ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660

agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag 780

tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt ggggaagacaa 900

Sequence Listing.ST25.txt

cctgtagggc	ctgcggggtc	tattgggaac	caagctggag	tgcagtggca	caatcttggc	960
tcactgcaat	ctccgcctcc	tgggttcaag	cgattctcct	gcctcagcct	cccgagttgt	1020
tgggattcca	ggcatgcatg	accaggctca	gctaattttt	gttttttttg	tagagacggg	1080
gtttcaccat	attggccagg	ctggtctcca	actcctaate	tcaggatgat	taccacactt	1140
ggcctcccaa	attgctggga	ttacaggcgt	gaaccactgc	tcccttccct	gtccttctga	1200
ttttaaaata	actataccag	caggaggacg	tccagacaca	gcataggcta	cctggccatg	1260
cccaaccggt	gggacatttg	agttgcttgc	ttggcactgt	cctctcatgc	gttggttcca	1320
ctcagtagat	gcctgttgaa	ttcgataccg	tcgacctcga	ggggggggccc	ggtaccagct	1380
tttgttccct	ttagtgaggg	ttaatttcga	gcttggcgta	atcatggtca	tagctgtttc	1440
ctgtgtgaaa	ttgttatccg	ctcacaattc	cacacaacat	acgagccgga	agcataaagt	1500
gtaaagcctg	gggtgcctaa	tgagtgaagt	aactcacatt	aattgcgttg	cgctcactgc	1560
ccgctttcca	gtcgggaaac	ctgtcgtgcc	agctgcatta	atgaatcggc	caacgcgcgg	1620
ggagaggcgg	tttgctgatt	gggcgtctt	ccgcttcctc	gctcactgac	tcgctgcgct	1680
cggtcgttcg	gctgcggcga	gcggtatcag	ctcactcaaa	ggcggttaata	cggttatcca	1740
cagaatcagg	ggataacgca	ggaaagaaca	tgtgagcaaa	aggccagcaa	aaggccagga	1800
accgtaaaaa	ggccgcgttg	ctggcgtttt	tccataggct	ccgccccct	gacgagcatc	1860
acaaaaatcg	acgctcaagt	cagaggtggc	gaaacccgac	aggactataa	agataccagg	1920
cgtttcccc	tggaagctcc	ctcgtgcgct	ctcctgttcc	gaccctgccg	cttaccggat	1980
acctgtccgc	ctttctccct	tcgggaagcg	tggcgctttc	tcatagctca	cgctgtaggt	2040
atctcagttc	ggtgtaggtc	gttcgctcca	agctgggctg	tgtgcacgaa	cccccggttc	2100
agcccgaccg	ctgcgcctta	tccggtaact	atcgtcttga	gtccaaccgc	gtaagacacg	2160
acttatcgcc	actggcagca	gccactggta	acaggattag	cagagcgagg	tatgtaggcg	2220
gtgctacaga	gttcttgaag	tgggtggccta	actacggcta	cactagaaga	acagtaattg	2280
gtatctgcgc	tctgctgaag	ccagttacct	tcggaaaaag	agttggtagc	tcttgatccg	2340
gcaaacaaac	caccgctggg	agcgggtggt	tttttggttg	caagcagcag	attacgcgca	2400
gaaaaaaagg	atctcaagaa	gatcctttga	tcttttctac	ggggtctgac	gtcagaaga	2460
actcgtcaag	aaggcgatag	aaggcgatgc	gctgcgaatc	gggagcggcg	ataccgtaaa	2520
gcacgaggaa	gcggtcagcc	cattcgccgc	caagctcttc	agcaatatca	cggttagcca	2580
acgctatgtc	ctgatagcgg	tccgccacac	ccagccggcc	acagtcgatg	aatccagaaa	2640
agcggccatt	ttccaccatg	atattcggca	agcaggcatc	gccatgggtc	acgacgagat	2700
cctcgccgtc	gggcatgcgc	gccttgagcc	tggcgaaacg	ttcggtggc	gcgagcccct	2760
gatgctcttc	gtccagatca	tcctgatcga	caagaccggc	ttccatccga	gtacgtgctc	2820

Sequence Listing.ST25.txt

```

gctcgatgcg atgtttcgct tgggtggcga atgggcaggt agccggatca agcgtatgca 2880
gccgccgcat tgcattcagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940
ggagatcctg ccccgccact tcgcccata gcagccagtc ctttcccgtc tcagtgacaa 3000
cgctcagcac agctgcgcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060
cgctcctgcag ttcatcagg gcaccggaca ggtcgggtctt gacaaaaaga accgggccc 3120
cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180
catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttggt 3240
caatcatgcy aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300
agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360
agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420
actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480
gatgtgctgc aaggcgatta agttgggtaa cgccagggtt tttccagtca cgac 3534

```

<210> 8
 <211> 3534
 <212> DNA
 <213> artificial sequence

<220>
 <223> This is the nucleic acid sequence for the operatively linked components of the TV-GHRH plasmid.

```

<400> 8
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60
accgcggtgg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120
gtgaggaatg gtggggagtt attttttagag cggtgaggaa ggtgggaggg cagcagggtgt 180
tggcgctcta aaaataactc ccgggagtta ttttttagagc ggaggaatgg tggacaccca 240
aatatggcga cggttcctca cccgtcgcca tatttggttg tccgccctcg gccggggccg 300
cattcctggg ggccggggcg tgctcccgcc cgctcagata aaaggctccg gggccggcg 360
cggcccacga gctaccggga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420
ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480
ctgggtgttc ttctttgtga tcctcacctc cagcaacagc tccactgct cccacctcc 540
ccctttgacc ctgaggatgc ggcggtatat cgatgccatc ttcaccaaca gctaccggaa 600
ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660
agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720
ggtggcatcc ctgtgacccc tcccagtgct ctctcctggc cctggaagtt gccactccag 780
tgcccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct 840

```

Sequence Listing.ST25.txt

tctataatat tatggggtgg aggggggtgg tatggagcaa ggggcaagtt ggggaagacaa	900
cctgtagggc ctgcgggggtc tattgggaac caagctggag tgcagtggca caatcttggc	960
tcactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt	1020
tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg	1080
gtttcaccat attggccagg ctggtctcca actcctaatac tcaggtgatc taccacacctt	1140
ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttcctt gtccttctga	1200
ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg	1260
cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca	1320
ctcagtagat gcctgttgaa ttcgataccg tcgacctcga gggggggccc ggtaccagct	1380
tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc	1440
ctgtgtgaaa ttgttatccg ctcaacaattc cacacaacat acgagccgga agcataaagt	1500
gtaaagcctg ggggtgcctaa tgagtgaagt aactcacatt aattgcgttg cgctcactgc	1560
ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg	1620
ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct	1680
cggtcgttcg gctgcggcga gcggtatcag ctactcaaaa ggcggtaata cggttatcca	1740
cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga	1800
accgtaaaaa ggccgcgttg ctggcgtttt tccataggct ccgccccctt gacgagcatc	1860
acaaaaatcg acgctcaagt cagagggtggc gaaacccgac aggactataa agataccagg	1920
cgtttcccc tggaaagctc ctctgctgct ctctgtttcc gacctgccg cttaccggat	1980
acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt	2040
atctcagttc ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa cccccgttc	2100
agcccgaacc ctgcgcctta tccggttaact atcgtcttga gtccaaccg gtaagacacg	2160
acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg	2220
gtgctacaga gttcttgaag tggtygccta actacggcta cactagaaga acagtatttg	2280
gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg	2340
gcaaacaaac caccgctggt agcggtyggt tttttgtttg caagcagcag attacgcgca	2400
gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gtcagaaga	2460
actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa	2520
gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca	2580
acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa	2640
agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat	2700

Sequence Listing.ST25.txt

```

cctcgccgctc gggcatgctc gccttgagcc tggcgaacag ttcggctggc gcgagcccct 2760
gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc 2820
gctcgatgctg atgtttcgct tgggtggcga atgggcaggt agccggatca agcgtatgca 2880
gccgccgcat tgcatacagc atgatggata ctttctcggc aggagcaagg tgagatgaca 2940
ggagatcctg ccccggcact tcgccaata gcagccagtc ctttcccgtc tcagtgacaa 3000
cgtcgagcac agctgctcaa ggaacgcccg tcgtggccag ccacgatagc cgcgctgcct 3060
cgtcctgcag ttcattcagg gcaccggaca ggctggctctt gacaaaaga accgggccc 3120
cctgcgctga cagccggaac acggcggcat cagagcagcc gattgtctgt tgtgcccagt 3180
catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 3240
caatcatgctg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc 3300
agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag 3360
agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca 3420
actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg 3480
gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac 3534

```

<210> 9
 <211> 3534
 <212> DNA
 <213> artificial sequence

<220>
 <223> This is the nucleic acid sequence for the operatively linked components of the 15/27/28 GHRH plasmid.

```

<400> 9
gttgtaaaac gacggccagt gaattgtaat acgactcact atagggcgaa ttggagctcc 60
accgcggttg cggccgtccg ccctcggcac catcctcacg acacccaaat atggcgacgg 120
gtgaggaatg gtggggagtt attttttagag cggtgaggaa ggtgggcagg cagcagggtgt 180
tggcgtctta aaaataactc ccgggagtta ttttttagagc ggaggaatgg tggacaccca 240
aatatggcga cggttcctca cccgtcgcca tatttgggtg tccgccctcg gccggggccg 300
cattcctggg ggccgggcgg tgctcccgcc cgcctcgata aaaggctccg gggccggcgg 360
cggcccacga gctacccgga ggagcgggag gcgccaagct ctagaactag tggatcccaa 420
ggcccaactc cccgaaccac tcagggtcct gtggacagct cacctagctg ccatggtgct 480
ctgggtgttc ttctttgtga tcctcaccct cagcaacagc tccactgct cccacctcc 540
ccctttgacc ctcaggatgc ggcggtatat cgatgccatc ttcaccaaca gctaccggaa 600
ggtgctggcc cagctgtccg cccgcaagct gctccaggac atcctgaaca ggcagcaggg 660
agagaggaac caagagcaag gagcataatg actgcaggaa ttcgatatca agcttatcgg 720

```


Sequence Listing.ST25.txt

ggtggcatcc ctgtgacccc tccccagtgc ctctcctggc cctggaagtt gccactccag	780
tgccccaccag ccttgtccta ataaaattaa gttgcatcat tttgtctgac taggtgtcct	840
tctataatat tatgggggtgg aggggggtgg tatggagcaa ggggcaagtt gggaagacaa	900
cctgtagggc ctgcggggtc tattgggaac caagctggag tgcagtggca caatcttggc	960
tactgcaat ctccgcctcc tgggttcaag cgattctcct gcctcagcct cccgagttgt	1020
tgggattcca ggcatgcatg accaggctca gctaattttt gtttttttgg tagagacggg	1080
gtttcaccat attggccagg ctggtctcca actcctaate tcaggtgatc taccacactt	1140
ggcctcccaa attgctggga ttacaggcgt gaaccactgc tcccttccct gtccttctga	1200
ttttaaaata actataccag caggaggacg tccagacaca gcataggcta cctggccatg	1260
cccaaccggt gggacatttg agttgcttgc ttggcactgt cctctcatgc gttgggtcca	1320
ctcagtagat gcctgttgaa ttcgataaccg tcgacctcga gggggggccc ggtaccagct	1380
tttgttccct ttagtgaggg ttaatttcga gcttggcgta atcatggtca tagctgtttc	1440
ctgtgtgaaa ttgttatccg ctcaaatc cacacaacat acgagccgga agcataaagt	1500
gtaaagcctg ggggtgcctaa tgagtgcctt aactcacatt aattgcgttg cgctcactgc	1560
ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg	1620
ggagaggcgg tttgcgtatt gggcgctctt ccgcttccct gctcactgac tcgctgcgct	1680
cggctggtcg gctgcggcga gcggtatcag ctactcaaa ggcggttaata cggttatcca	1740
cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga	1800
accgtaaaaa ggcgcggttg ctggcggttt tccataggct ccgccccctt gacgagcatc	1860
acaaaaatcg acgctcaagt cagagggtggc gaaacccgac aggactataa agataaccagg	1920
cgtttcccc tggaaagctcc ctctgtgcgt ctctgttcc gaccctgccg cttaccggat	1980
acctgtccgc ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtaggt	2040
atctcagttc ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa cccccgttc	2100
agcccgaccg ctgcgctta tccggttaact atcgtcttga gtccaaccg gtaagacacg	2160
acttatcgcc actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg	2220
gtgctacaga gttcttgaag tgggtggccta actacggcta cactagaaga acagtatttg	2280
gtatctgcgc tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg	2340
gcaaacaaac caccgtggt agcgggtggtt tttttgtttg caagcagcag attacgcgca	2400
gaaaaaaagg atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagaaga	2460
actcgtcaag aaggcgatag aaggcgatgc gctgcgaatc gggagcggcg ataccgtaaa	2520
gcacgaggaa gcggtcagcc cattcgccgc caagctcttc agcaatatca cgggtagcca	2580
acgctatgtc ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa	2640

Sequence Listing.ST25.txt

agcggccatt ttccaccatg atattcggca agcaggcatc gccatgggtc acgacgagat	2700
cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag ttcggctggc gcgagcccc	2760
gatgctcttc gtccagatca tcctgatcga caagaccggc ttccatccga gtacgtgctc	2820
gctcgatgcg atgtttcgtc tgggtggcga atgggcaggc agccggatca agcgtatgca	2880
gccgccgcat tgcacagacc atgatggata ctttctcggc aggagcaagg tgagatgaca	2940
ggagatcctg ccccggcact tcgccaata gcagccagtc ctttcccgtc tcagtgacaa	3000
cgtcgagcac agctgcgcaa ggaacgcccgc tcgtggccag ccacgatagc cgcgctgcct	3060
cgtcctgcag ttcatcagc gcaccggaca ggtcggctct gacaaaaaga accggggcgc	3120
cctgcgctga cagccggaac acggcgccat cagagcagcc gattgtctgt tgtgcccagt	3180
catagccgaa tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttggt	3240
caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc ctgcgccatc	3300
agatccttgg cggcaagaaa gccatccagt ttactttgca gggcttccca accttaccag	3360
agggcgcccc agctggcaat tccggttcgc ttgctgtcca taaaaccgcc cagtctagca	3420
actgttggga agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg	3480
gatgtgctgc aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgac	3534